

Sharp Constant in Jackson's Inequality with Modulus of Smoothness for Uniform Approximations of Periodic Functions

S. A. Pichugov*

Dnepropetrovsk National Technical University of Railroad Communications

Received April 22, 2012

Abstract—It is proved that, in the space $C_{2\pi}$, for all $k, n \in \mathbb{N}, n > 1$, the following inequalities hold:

$$\left(1 - \frac{1}{2n}\right) \frac{k^2 + 1}{2} \leq \sup_{\substack{f \in C_{2\pi} \\ f \neq \text{const}}} \frac{e_{n-1}(f)}{\omega_2(f, \pi/(2nk))} \leq \frac{k^2 + 1}{2}.$$

where $e_{n-1}(f)$ is the value of the best approximation of f by trigonometric polynomials and $\omega_2(f, h)$ is the modulus of smoothness of f . A similar result is also obtained for approximation by continuous polygonal lines with equidistant nodes.

DOI: 10.1134/S0001434613050118

Keywords: *Jackson's inequality, periodic function, trigonometric polynomial, modulus of smoothness, polygonal line, Steklov mean, Favard sum.*

Suppose that

- $C_{2\pi}$ is the space of (2π) -periodic real-valued continuous functions f with norm

$$\|f\| = \max\{|f(x)| : x \in \mathbb{R}\};$$

- $e_{n-1}(f) = \inf_{T_{n-1}} \|f - T_{n-1}\|$ is the value of the best approximation of f in this space by trigonometric polynomials T_{n-1} of degree at most $n - 1$, $n \in \mathbb{N}$;
- $\omega_2(f, h) = \sup_{|t| \leq h} \|\Delta_t^2 f\|$ is the value of the modulus of smoothness of f at a point h , $h \geq 0$, where

$$\Delta_t^2 f(x) = f(x + t) + f(x - t) - 2f(x)$$

is the second difference of f at a point x with step t .

Theorem 1. *For all $k, n \in \mathbb{N}, n > 1$, the following inequalities hold:*

$$\left(1 - \frac{1}{2n}\right) \frac{k^2 + 1}{2} \leq \sup_{\substack{f \in C_{2\pi} \\ f \neq \text{const}}} \frac{e_{n-1}(f)}{\omega_2(f, \pi/(2nk))} \leq \frac{k^2 + 1}{2}. \quad (1)$$

Corollary 1. *For all $k \in \mathbb{N}$, the following relations hold:*

$$\sup_n \sup_{\substack{f \in C_{2\pi} \\ f \neq \text{const}}} \frac{e_{n-1}(f)}{\omega_2(f, \pi/(2nk))} = \frac{k^2 + 1}{2}.$$

*E-mail: s.a.pichugov@mail.ru

Upper bounds for the values of best approximations of functions in terms of the values of their moduli of continuity of various orders are called the *Jackson's inequalities*. Well-known results concerning sharp Jackson inequalities (i.e., inequalities with sharp constants) for functions of one variable can be found in [1]–[8]. In particular, in the case $k = 1$, inequalities (1) were proved in [9], [10] (upper bound) and [6] (lower bound). Also note the paper [11] in which, for other values of the argument of the modulus of smoothness, upper bounds for sharp constants were obtained.

Suppose that M is an arbitrary subspace in $C_{2\pi}$ containing constants,

$$e(f; M) = \inf\{\|f - g\| : g \in M\}$$

is the value of the best approximation of f by the subspace M ,

$$\mathcal{W}^2 = \{f \in C_{2\pi} : f' \in AC, f'' \in C_{2\pi}, \|f''\| \leq 1\},$$

and $e(\mathcal{W}^2; M)$ is the value of the best approximation of the class \mathcal{W}^2 by the subspace M .

Lemma 1. 1) For any f from $C_{2\pi}$, the following inequality holds:

$$e(f; M) \leq \frac{1}{2} \inf_{h>0} \left(1 + \frac{2e(\mathcal{W}^2; M)}{h^2} \right) \omega(f, h); \quad (2) \quad \{\text{eq2:v932}\}$$

2) for any $\delta > 0$, the following inequalities hold:

$$\frac{\delta^2}{2} \leq \sup_{\substack{f \in C_{2\pi} \\ f \neq \text{const}}} \frac{e(f; M)}{\omega_2(f, (2e(\mathcal{W}^2; M))^{1/2}/\delta)} \leq \frac{\delta^2 + 1}{2}. \quad (3) \quad \{\text{eq3:v932}\}$$

Proof of Lemma 1. For $h > 0$, suppose that

$$\mathcal{S}_h(f, x) = \frac{1}{h} \int_{-h/2}^{h/2} f(x+t) dt$$

is the Steklov mean of f with step h , and

$$\mathcal{S}_{h^2}(f, x) := \mathcal{S}_h(\mathcal{S}_h f, x) = \frac{1}{h^2} \int_{-h}^h (h - |t|) f(x+t) dt$$

is the Steklov mean of f of second order. Then

$$\begin{aligned} |f(x) - \mathcal{S}_{h^2}(f, x)| &\leq \frac{1}{h^2} \int_0^h (h-t) |\Delta_t^2 f(x)| dt, \\ \|f - \mathcal{S}_{h^2} f\| &\leq \frac{1}{h^2} \int_0^h (h-t) \omega_2(f, t) dt \leq \frac{1}{2} \omega_2(f, h). \end{aligned}$$

Further,

$$\|D^2(\mathcal{S}_{h^2} f)\| = \left\| \frac{\Delta_h^2 f}{h^2} \right\| \leq \frac{\omega_2(f, h)}{h^2}, \quad e(\mathcal{S}_{h^2} f; M) \leq \frac{1}{h^2} \omega_2(f, h) e(\mathcal{W}^2; M).$$

Now, to find an upper bound for the approximation value $e(f; M)$, we use the intermediate approximation of f by smoother functions $\mathcal{S}_{h^2} f$:

$$e(f; M) \leq \|f - \mathcal{S}_{h^2} f\| + e(\mathcal{S}_{h^2} f; M) \leq \frac{1}{2} \left(1 + \frac{2}{h^2} e(\mathcal{W}^2; M) \right) \omega_2(f, h). \quad (4) \quad \{\text{eq4:v932}\}$$

Since the value of h is arbitrary, we obtain (2). Note that this method was used in [9] to find estimate (4) for the approximation by polynomials.

If we put $h = (2e(\mathcal{W}^2; M))^{1/2}/\delta$ in (4), then we obtain the upper bound in (3). We obtain the lower bound in (3) by restricting ourselves to the approximation of smooth functions f from $C_{2\pi}$ and using the inequality $\omega_2(f, h) \leq \|f''\| h^2$.

Lemma 1 is proved. \square

Proof of Theorem 1. In the case of approximation by trigonometric polynomials using the Akhiezer–Krein–Favard theorem (see, for example, [7]), we obtain

$$\sup_{f \in \mathcal{W}^2} e_{n-1}(f) = \frac{\pi^2}{8n^2}, \quad (5) \quad \{\text{eq5:v932}\}$$

and then the upper bound in (3) is of the form

$$\sup_{\substack{f \in C_{2\pi} \\ f \neq \text{const}}} \frac{e_{n-1}(f)}{\omega_2(f, \pi/(2n\delta))} \leq \frac{\delta^2 + 1}{2}. \quad (6) \quad \{\text{eq6:v932}\}$$

Let us show that, for $\delta \in \mathbb{N}$, this estimate cannot be improved for all n .

To find lower bounds for the Jackson constants in the construction of the following functions we use an idea of Korneichuk [1], [2], which was realized in [6] for the moduli of smoothness for $\delta = 1$.

Let us fix

$$k, n \in \mathbb{N}, \quad n > 1, \quad \varepsilon \in \left(0, \frac{1}{2}\right],$$

and set

$$x_0 = 0, \quad x_\nu = \nu h - (n - \nu)\beta, \quad \nu = 1, \dots, n, \quad h = \frac{\pi}{n}, \quad \beta \in \left(0, \frac{4\varepsilon}{n^2(k^2 + 1)}\right).$$

By construction,

$$x_{\nu+1} - x_\nu = h + \beta, \quad x_n = \pi.$$

Consider an arbitrary function f from $C_{2\pi}$ satisfying the conditions

$$f(-x) = f(x), \quad f(0) = 0, \quad f(x_\nu) = (-1)^{\nu+1} \frac{k^2 + 1}{2}, \quad \nu = 1, \dots, n. \quad (7) \quad \{\text{eq7:v932}\}$$

To find a lower bound for $e_{n-1}(f)$, we use the polynomial

$$T_{n-1}(x) = \frac{k^2 + 1}{2n} \frac{\sin(n - 1/2)x}{2 \sin(x/2)}.$$

For $\nu = 0, 1, \dots, n$, we have (see [1], [2])

$$f(x_\nu) - T_{n-1}(x_\nu) = (-1)^{\nu+1} \left(\frac{k^2 + 1}{2} - \frac{k^2 + 1}{4n} \right) + \mu_\nu,$$

where $|\mu_\nu| < \varepsilon$; hence, taking into account the fact that f is even and using the Vallée–Poussin theorem, we obtain

$$e_{n-1}(f) \geq \frac{k^2 + 1}{2} \left(1 - \frac{1}{2n} \right) - \varepsilon. \quad (8) \quad \{\text{eq8:v932}\}$$

Let us now define the function $f(x)$ on the whole axis so that, along with conditions (7), the following condition also holds:

$$\omega_2\left(f, \frac{\pi}{2nk}\right) = 1. \quad (9) \quad \{\text{eq9:v932}\}$$

First, let us construct $f(x)$ on the closed interval $[x_1, \gamma]$, where $\gamma = (3/2)(h + \beta) - n\beta$ is the midpoint of the closed interval $[x_1, x_2]$, specifying it the polygonal line uniquely defined by its values at the nodes:

$$\begin{aligned} f(\gamma) &= 0, \quad f\left(x_1 + j \frac{h}{2k}\right) = \frac{k^2 + 1}{2} - \frac{j^2}{2}, \quad j = 0, \dots, k, \\ f\left(x_1 + j \frac{h}{2k} + \frac{\beta}{2}\right) &= \frac{k^2 + 1}{2} - \frac{(j+1)^2}{2}, \quad j = 0, \dots, k-1. \end{aligned} \quad (10) \quad \{\text{eq10:v932}\}$$

Let us continue $f(x)$ to the closed interval $[\gamma, x_2]$ as an odd function with respect to the point γ :

$$f(\gamma + x) = -f(\gamma - x), \quad x \in \left[0, \frac{x_2 - x_1}{2}\right]. \quad (11) \quad \{\text{eq11:v93}\}$$

Further, we set

$$\begin{aligned} f(x) &= -f(x - h - \beta), & x &\in [x_2, \pi], \\ f(x) &= \max\{0; f(2x_1 - x)\}, & x &\in [0, x_1], \\ f(-x) &= f(x), & x &\in [-\pi; 0], \\ f(x + 2\pi) &= f(x). \end{aligned} \quad (12) \quad \{\text{eq12:v93}\}$$

This defines the continuous 2π -periodic function satisfying conditions (7). It is easy to see that condition (9) also holds: since $f(x)$ is a polygonal line, it follows that, to calculate its modulus of smoothness, it suffices to calculate the increments of the function f at its nodes.

Since ε is arbitrary, relations (8) and (9) imply the lower bound of the Jackson constant in (1).

Theorem 1 is proved. \square

Remark 1. In the proof of Lemmas 1, we did not use the specific properties of the metric of $C_{2\pi}$; in particular, relations (3) are also valid in the space $L_1[0, 2\pi]$. Further, the analog (5) of the Akhiezer–Krein–Favard Theorem also holds in $L_1[0, 2\pi]$ (see, for example, [7]). Therefore, in the space $L_1[0, 2\pi]$, the following upper bound similar to (6) is also valid:

$$\sup_{\substack{f \in L_1[0, 2\pi] \\ f \neq \text{const}}} \frac{e_{n-1}(f)_{L_1}}{\omega_2(f, \pi/(2n\delta))_{L_1}} \leq \frac{\delta^2 + 1}{2}.$$

However, we do not know the exact values of the Jackson constants for the moduli of smoothness in this space for any $\delta > 0$.

Remark 2. Suppose that $X_{n-1,2}(f)$ are the Favard sums of degree $n - 1$ of order 2 for the function f (see, for example, [7]). Then

$$\widetilde{\mathcal{L}}_{n-1}(f) := \mathcal{S}_{(\pi/(2nk))^2} \circ X_{n-1,2}(f)$$

is the best linear method for approximating functions among all linear polynomial methods \mathcal{L}_{n-1} in the sense that, for any $k \in \mathbb{N}$,

$$\sup_n \inf_{\mathcal{L}_{n-1}} \sup_{\substack{f \in C_{2\pi} \\ f \neq \text{const}}} \frac{\|f - \mathcal{L}_{n-1}(f)\|}{\omega_2(f, \pi/(2nk))} = \sup_n \sup_{\substack{f \in C_{2\pi} \\ f \neq \text{const}}} \frac{\|f - \widetilde{\mathcal{L}}_{n-1}(f)\|}{\omega_2(f, \pi/(2nk))} = \frac{k^2 + 1}{2}.$$

This immediately follows from the proof of the upper bound in Theorem 1 and the fact (see, for example, [7]) that

$$e_{n-1}(\mathcal{W}^2) = \sup_{f \in \mathcal{W}^2} \|f - X_{n-1,2}(f)\|.$$

It is easy to calculate the multipliers of the method $\widetilde{\mathcal{L}}_{n-1}$: if f_ν are the complex Fourier coefficients of f and

$$\widetilde{\mathcal{L}}_{n-1}(f, x) = \sum_{|\nu| < n} \alpha_k\left(\frac{\nu}{n}\right) f_\nu e^{i\nu x},$$

then

$$\alpha_k(t) = 4k^2 \left(\sin \frac{\pi}{4k} t \right)^2 \frac{\cos(\pi t/2)}{(\sin(\pi t/2))^2}, \quad |t| \leq 1.$$

In particular (see [9], [10], [6]),

$$\alpha_1(t) = 1 - \left(\tan \frac{\pi}{4} t \right)^2, \quad \alpha_2(t) = \frac{1 - (\tan(\pi t/4))^2}{(\cos(\pi t/8))^2}.$$

Let us also consider the approximation of functions by the subspace \mathcal{S}_{2n} of periodic continuous polygonal lines with the $2n$ equidistant nodes

$$y_\nu = \frac{\pi}{2n} + \frac{\nu\pi}{n}, \quad \nu \in \mathbb{Z},$$

on the period $[-\pi, \pi]$.

Theorem 2. *For all $k, n \in \mathbb{N}, n > 1$, the following inequalities hold:*

$$\left(1 - \frac{1}{2n}\right) \frac{k^2 + 1}{2} \leq \sup_{\substack{f \in \mathcal{C}_{2\pi} \\ f \neq \text{const}}} \frac{e(f; \mathcal{S}_{2n})}{\omega_2(f, \pi/(2nk))} \leq \frac{k^2 + 1}{2}. \quad (13)$$

Corollary 2. *For all $k \in \mathbb{N}$, the following relations hold:*

$$\sup_n \sup_{\substack{f \in \mathcal{C}_{2\pi} \\ f \neq \text{const}}} \frac{e(f; \mathcal{S}_{2n})}{\omega_2(f, \pi/(2nk))} = \frac{k^2 + 1}{2}.$$

Proof. Since (see [11])

$$e(\mathcal{W}^2; \mathcal{S}_{2n}) = \frac{\pi^2}{8n^2},$$

we see that the upper bound in (13) follows from (3).

To find the lower bound, we consider the approximation of the function f constructed in the proof of Theorem 1 (see (7), (10)–(12)). We shall use the duality relation for approximation by splines of minimal deficiency [12]; in our particular case, this relation can be expressed as

$$e(f; \mathcal{S}_{2n}) = \sup \left\{ \int_{-\pi}^{\pi} f(x) dg_1(x) : \text{Var } g_1(x) \leq 1, g_2(y_\nu) = \text{const}, \nu \in \mathbb{Z} \right\}, \quad (14)$$

where $g_2(x)$ is the antiderivative of $g_1(x)$, which is zero in the mean, and $\text{Var } g_1(x)$ is the variation of $g_1(x)$ on the period.

To find the lower bound for $e(f; \mathcal{S}_{2n})$, we construct a piecewise constant function $g_1(x)$ as follows: first, we define the auxiliary function $\psi(x)$ on the period $[-\pi, \pi]$ as an even continuous polygonal line with zeros at the points y_ν and the vertices at the points x_ν .

For $x \in [0, \pi]$, let

$$\psi(x) := c_\nu(x - y_\nu), \quad x \in [x_{\nu-1}, x_\nu], \quad \nu = 1, \dots, n.$$

The continuity condition at the point $x_{\nu+1}$ means that

$$c_{\nu+1} = -c_\nu \frac{\pi/(2n) - (n - (\nu + 1))\beta}{\pi/(2n) + (n - (\nu + 1))\beta}.$$

Set $c_1 = -1$; then, for $\nu = 2, \dots, n$,

$$c_\nu = (-1)^\nu \prod_{j=1}^{\nu-1} \frac{\pi/(2n) - (n - j)\beta}{\pi/(2n) + (n - j)\beta}. \quad (15)$$

The function $\psi'(x)$ is piecewise constant and

$$\text{Var } \psi'(x) = 4 \sum_{\nu=1}^n |c_\nu|.$$

Set

$$g_2(x) = \frac{\psi(x) - \psi_0}{4 \sum_{\nu=1}^n |c_\nu|}, \quad \text{where } \psi_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} \psi(x) dx.$$

Then $g_2(x)$ is zero in the mean, $g_2(y_\nu) = \text{const}$, $\nu \in \mathbb{Z}$, and $\text{Var } g_1(x) = 1$. Since

$$|c_\nu - c_{\nu+1}| = |c_\nu| + |c_{\nu+1}|,$$

(see (15)), it follows from (14) that

$$\begin{aligned} e(f; \mathcal{S}_{2n}) &\geq \int_{-\pi}^{\pi} f(x) dg_1(x) = \frac{1}{4 \sum_{\nu=1}^n |c_\nu|} 2 \left(\sum_{\nu=1}^{n-1} |c_\nu - c_{\nu+1}| + |c_n| \right) \frac{k^2 + 1}{2} \\ &= \frac{1}{4 \sum_{\nu=1}^n |c_\nu|} \left(4 \sum_{\nu=1}^n |c_\nu| - 2|c_1| \right) \frac{k^2 + 1}{2} = \left(1 - \frac{1}{2 \sum_{\nu=1}^n |c_\nu|} \right) \frac{k^2 + 1}{2}. \end{aligned}$$

Equality (15) implies that $|c_\nu| \rightarrow 1$ as $\beta \rightarrow 0$. This yields the lower bound in (13). Theorem 2 is proved. \square

It is possible that the assertion of Theorem 2 remains valid in the case of approximation by splines of minimal deficiency and any order $r \in \mathbb{N}$. In this case, the upper bound in (13) holds and it suffices only to prove the lower bound.

REFERENCES

1. N. P. Korneichuk, "The exact constant in Jackson's theorem on best uniform approximation of continuous periodic functions," *Dokl. Akad. Nauk SSSR* **145**(3), 514–515 (1962) [*Soviet Math. Dokl.* **3**(3), 1040–1041 (1962)].
2. N. P. Korneichuk, "On the sharp constant in Jackson's inequality for continuous periodic functions," *Mat. Zametki* **32**(5), 669–674 (1982) [*Math. Notes* **32**(5), 818–821 (1983)].
3. N. I. Chernykh, "On Jackson's inequality in L_2 ," in *Trudy Mat. Inst. Steklov*, Vol. 88: *Approximation of Functions in the Mean*, Collection of papers (Nauka, Moscow, 1967), pp. 71–74 [*Proc. Steklov Inst. Math.* **88**, 75–78 (1967)].
4. N. I. Chernykh, "Best approximation of periodic functions by trigonometric polynomials in L_2 ," *Mat. Zametki* **2**(5), 513–522 (1967) [*Math. Notes* **2**(5), 803–808 (1968)].
5. N. I. Chernykh, "Jackson's inequality in $L_p(0, 2\pi)$, ($1 \leq p < 2$), with sharp constant," in *Trudy Mat. Inst. Steklov*, Vol. 198: *Proceedings of an All-Union School on the Theory of Functions, Miass, July 1989* (Nauka, Moscow, 1992), pp. 232–241 [*Proc. Steklov Inst. Math.* **198**, 223–231 (1994)].
6. V. V. Shalaev "On the approximation of continuous periodic functions by trigonometric polynomials," in *Studies of Problems of Current Interest Dealing with Summation and Approximations of Functions and Their Applications* (Dnepropetrovsk. Univ., Dnepropetrovsk, 1979), pp. 39–43 [in Russian].
7. N. P. Korneichuk, *Exact Constants in Approximation Theory* (Nauka, Moscow, 1987) [in Russian].
8. A. A. Ligon, "Sharp constants in inequalities of Jackson type," in *Special Questions of Approximation Theory and Optimal Control of Distributed Systems* (Vishcha Shkola, Kiev, 1990), pp. 3–74 [in Russian].
9. V. V. Zhuk, "Some sharp inequalities between best approximations and moduli of continuity," *Vestnik Leningrad. Univ. Mat. Mekh. Astronom.*, No. 1, 21–26 (1974).
10. V. V. Zhuk, *Approximation of Periodic Functions* (Izd. Leningradsk. Univ., Leningrad, 1982) [in Russian].
11. A. G. Babenko and Yu. V. Kryakin, "Integral approximation of the characteristic function of the interval and Jackson's inequality in $C(\mathbb{T})$," *Trudy Inst. Mat. Mekh. (Ural Branch, Russian Academy of Sciences)* (2009), Vol. 15, pp. 59–65 [in Russian].
12. N. P. Korneichuk, *Splines in Approximation Theory* (Nauka, Moscow, 1984) [in Russian].